

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
**WO 03/013224 A2**

(51) International Patent Classification<sup>7</sup>: **A01H**

(21) International Application Number: PCT/EP02/08136

(22) International Filing Date: 19 July 2002 (19.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/921,922 6 August 2001 (06.08.2001) US

(71) Applicant: **BAYER BIOSCIENCE N.V.** [BE/BE]; Jozef Plateaustraat 22, B-9000 Gent (BE).

(72) Inventors: **TROLINDER, Linda**; 1004 Dogwood St, Idalout, TX 79329 (US). **JEFFERSON, Gwyn**; 2787 Marico Place, Green Ville, MS 38701 (US). **DE BEUCK-  
ELEER, Marc**; Hekers 36, B-9052 Zwijnaarde (BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,

MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**


- as to the identity of the inventor (Rule 4.17(i)) for all designations
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HERBICIDE TOLERANT COTTON PLANTS AND METHODS FOR PRODUCING AND IDENTIFYING SAME



(57) Abstract: The invention pertains to transgenic cotton plants, plant material and seeds, characterized by harboring a specific transformation event, particularly by the presence of a gene encoding a protein that confers herbicide tolerance, at a specific location in the cotton genome. The cotton plants of the invention combine the herbicide tolerant phenotype with optimal agronomic performance.



**WO 03/013224 A2**

Title of the invention**HERBICIDE TOLERANT COTTON PLANTS AND METHODS FOR  
PRODUCING AND IDENTIFYING SAME**

5

Field of the invention

This invention pertains to transgenic cotton plants, plant material and seeds, characterized by harboring a specific transformation event, particularly by the presence of a gene  
10 encoding a protein that confers herbicide tolerance, at a specific location in the cotton genome. The cotton plants of the invention combine the herbicide tolerant phenotype with an agronomic performance, genetic stability and adaptability to different genetic backgrounds equivalent to the non-transformed cotton line in the absence of weed pressure.

15

All documents cited herein are hereby incorporated herein by reference.

Background of the invention

20 The phenotypic expression of a transgene in a plant is determined both by the structure of the gene itself and by its location in the plant genome. At the same time the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant in different ways. The agronomically or industrially successful introduction of a commercially interesting trait in a plant by genetic manipulation can be a lengthy  
25 procedure dependent on different factors. The actual transformation and regeneration of genetically transformed plants are only the first in a series of selection steps, which include extensive genetic characterization, breeding, and evaluation in field trials.

30 Cotton fiber is the single most important textile worldwide. About 80 million acres of cotton are harvested annually across the globe. Cotton is the fifth largest crop in the U.S. in terms of acreage production, with over 15 million acres planted in 2000. Primary weed

species for cotton are *Ipomoea* sp. (morning glory), *Amaranthus* spp. (pigweed), *Cyperus* spp. (nutsedge), *Xanthium* spp. (cocklebur) and *Sorghum* spp. (johnsongrass). Before the introduction of broad-leaf herbicides that could be used on a growing cotton field, growers used directed, post-emergence applications of nonselective herbicides taking  
5 care not to contact the growing crop plants. As this requires a difference in height between the weeds and the crop, this is not always possible. Especially for small cotton, this practice is time-consuming and potentially damaging to the crop.

The *bar* gene (Thompson et al, 1987, EMBO J 6:2519-2523; Deblock *et al.* 1987, EMBO  
10 J. 6:2513-2518) is a gene encoding the enzyme phosphinothricin acetyl transferase (PAT), which, when expressed in a plant, confers resistance to the herbicidal compounds phosphinothricin (also called glufosinate) or bialaphos (see also for example US patents, 5,646,024 and 5,561,236) and salts and optical isomers thereof. Phosphinothricin controls broadleaf weeds including morning glory and has a wide window of application.

15 Successful genetic transformation of cotton has been obtained by a number of methods including *Agrobacterium* infection of cotton explants (Firoozabady et al. 1987, Plant Molecular Biology 10:105-116; Umbeck et al. 1987, Bio/Technology 5:263-266 and in WO 00/71733, US 5,004,863, and US 5,159,135), as well as direct gene transfer by  
20 microprojectile bombardment of meristematic cotton tissues (Finer and Mc Mullen, 1990, Plant Cell Reports, 5:586-589; McCabe and Martinell, 1993, Bio/Technology 11:596-598, WO92/15675, EP0 531 506). Increased transformation efficiency for *Agrobacterium* transformation has been reported using the methods described by Hansen et al. (1994, Proc. Nat. Acad. Sci. 91:7603-7607) Veluthambi et al. (1989, Journal of Bacteriology  
25 171:3696-3703) and WO 00/71733.

Different methods for regeneration of cotton plants have also been described (WO 89/05344, US 5,244,802, US 5,583,036, WO89/12102, WO98/15622, and WO97/12512).

However, the foregoing documents fail to teach or suggest the present invention.

### Summary of the invention

The present invention relates to a transgenic cotton plant, or seed, cells or tissues thereof, comprising, stably integrated into its genome, an expression cassette which comprises a herbicide tolerance gene comprising the coding sequence of the *bar* gene (as described in Example 1.1 herein), which is herbicide tolerant and, in the absence of weed pressure, has an agronomic performance which is substantially equivalent to the non-transgenic isolate. Under weed pressure and the appropriate Liberty<sup>TM</sup> treatment, the plant will have a superior agronomic phenotype compared to the non-transgenic plant.

In one embodiment of the invention, the cotton plant or seed, cells or tissues thereof, comprises the expression cassette of pGSV71 (as described in Example 1.1, Table 1 herein). In the preferred embodiment of the invention the cotton plant or seed, cells or tissues thereof comprise elite event EE-GH1.

In another embodiment of the invention, the transgenic cotton plant or seed, cells or tissues thereof comprises:

- (i) event EE-GH1 in its genome ;or
- (ii) event EE-GH1 with the proviso that the *bar* gene used in the event is substituted with a nucleic acid sequence that hybridizes to the complement of the *bar* gene under stringent conditions.

More specifically, the present invention relates to a transgenic cotton plant, seed, cells or tissues thereof, the genomic DNA of which is characterized by the fact that, when analyzed in a PCR identification protocol as described herein, using two primers directed to the 5' or 3' flanking region of EE-GH1 and the foreign DNA, respectively, yields a fragment which is specific for EE-GH1. Preferably the primers are directed against the 5' flanking region within SEQ ID NO: 1 and the foreign DNA respectively; most preferably, the primers comprise the nucleotide sequence of SEQ ID NO: 2 and SEQ ID NO: 3 respectively, and yield a DNA fragment of between 250 and 290 bp, preferably of about 269 bp.

Reference seed comprising the elite event of the invention has been deposited at the ATCC under accession number PTA-3343. Thus, a preferred embodiment of the invention is the seed comprising elite event EE-GH1 deposited as ATCC accession  
5 number PTA-3343, which will grow into a cotton plant resistant to glufosinate. The seed of ATCC deposit number PTA-3343, which is a seed lot consisting of about 50% non-transgenic kernels and 50% transgenic kernels hemizygous for the transgene, comprising the elite event of the invention, which will grow into glufosinate tolerant plants. The seed can be sown and the growing plants can be treated with PPT or Liberty<sup>TM</sup> as described  
10 herein to obtain 100% glufosinate tolerant plants, comprising the elite event of the invention. The invention further relates to cells, tissues, progeny, and descendants from a plant comprising the elite event of the invention grown from the seed deposited at the ATCC having accession number PTA-3343. The invention further relates to plants obtainable by propagation of and/or breeding with a cotton plant comprising the elite  
15 event of the invention grown from the seed deposited at the ATCC having accession number PTA-3343.

The invention further relates to plants, seeds, cells or tissues comprising a foreign DNA sequence, preferably a herbicide tolerance gene as described herein, integrated into the  
20 chromosomal DNA in a region which comprises the plant DNA sequence of SEQ ID NO: 1 and/or SEQ ID NO: 4, more particularly which comprises the DNA sequence of SEQ ID NO: 5, or a sequence which hybridizes under stringent conditions to a sequence that is complementary to a sequence comprising the plant DNA sequence of SEQ ID NO: 1, SEQ ID NO: 4 and/or SEQ ID NO: 5.

25 The invention further provides a process for producing a transgenic cell of a cotton plant, which comprises inserting a recombinant DNA molecule into a region of the chromosomal DNA of a cotton cell, tissue or callus which comprises the plant DNA sequence of SEQ ID NO: 1 and/or SEQ ID NO: 4, or which comprises a sequence which  
30 hybridizes under stringent conditions to a sequence that is complementary to a sequence comprising the plant DNA sequence of SEQ ID NO: 1 and/or SEQ ID NO: 4.

The invention further relates to a method for identifying a transgenic plant, or cells or tissues thereof, comprising elite event EE-GH1 which method is based on identifying the presence of characterizing DNA sequences or amino acids encoded by such DNA sequences in the transgenic plant, cells or tissues. According to a preferred embodiment of the invention, such characterizing DNA sequences are sequences of 15bp, preferably 20bp, most preferably 30bp or more which comprise the insertion site of the event, i.e. both a part foreign DNA and a part of the cotton genome (either the 5' or 3' flanking region) contiguous therewith, allowing specific identification of the elite event.

According to another aspect of the invention, DNA sequences are disclosed comprising the insertion site of the event and sufficient length of polynucleotides of both the cotton genomic DNA and the foreign DNA (transgene), so as to be useful as primer or probe for the detection of EE-GH1. Preferably, such sequences comprise at least 9 nucleotides of the cotton genomic DNA and a similar number of nucleotides of the foreign DNA (transgene) of EE-GH1 therewith at each side of the insertion site respectively. Most preferably, such DNA sequences comprise at least 9 nucleotides of the cotton genomic DNA and a similar number of nucleotides of the foreign DNA contiguous with the insertion site in SEQ ID NO: 1 or SEQ ID NO: 4.

According to another preferred aspect of the invention, the method for identifying a transgenic plant, or cells or tissues thereof, comprising elite event EE-GH1, comprises amplifying a sequence of a nucleic acid present in biological samples, using a polymerase chain reaction, with at least two primers, one of which recognizes the plant DNA in the 5' or 3' flanking region of EE-GH1, the other which recognizes a sequence within the foreign DNA. Preferably, the genomic DNA is analyzed using primers which recognize a sequence within the plant 5' flanking region of EE-GH1, most preferably within the plant DNA sequence in SEQ ID NO: 1, and a sequence within the foreign DNA, respectively. Especially preferably, the genomic DNA is analyzed according to the PCR identification protocol described herein whereby the primer recognizing a sequence within the 5' flanking region comprises the nucleotide sequence of SEQ ID NO: 2. Particularly, the

primer recognizing a sequence within the 5' flanking region comprises the nucleotide sequence of SEQ ID NO: 2 and the primer recognizing a sequence within the foreign DNA comprises the nucleotide sequence of SEQ ID NO: 3, so that the amplified fragment is a fragment preferably of between 250 and 290 bp, more preferably of  
5 between 260 and 270 bp, most preferably of about 269 bp.

Accordingly, the present invention relates to the transgenic plant, cells or tissues thereof which can be identified according the above-described identification methods for EE-GH1.

10 The present invention relates to methods for identifying elite event EE-GH1 in biological samples, which methods are based on primers or probes that specifically recognize the 5' and/or 3' flanking sequence of EE-GH1. In a preferred embodiment of the invention these methods are based on primers or probes which recognize a sequence within SEQ ID  
15 NO: 1 and/or SEQ ID NO: 4, more particularly primers or probes comprising the sequence of SEQ ID NO: 2. According to another embodiment of the invention such methods are based on primers or probes which recognize a characterizing sequence of EE-GH1, which is a DNA sequence comprising the insertion site of the event, including sufficient nucleotides of the 5' or 3' flanking DNA and the foreign DNA contiguous  
20 therewith so as to be diagnostic for the event. Most particularly, such methods are based on sequences comprising sufficient nucleotides of the 5' or 3' flanking DNA and the foreign DNA contiguous therewith of SEQ ID NO: 1 or SEQ ID NO: 4. Such methods are known in the art and include, but are not limited to, the method described by Lyamichev et al. (1999, Nature Biotechnol 17:292-296).

25 The present invention further relates to the specific flanking sequences of EE-GH1 described herein, which can be used to develop specific identification methods for EE-GH1 in biological samples. More particularly, the invention relates to the 5' and or 3' flanking regions of EE-GH1, which can be used for the development of specific primers  
30 and probes as well as to the specific primers and probes developed from the 5' and/or 3' flanking sequences of EE-GH1 and the insertion site adjacent thereto. The invention

further relates to identification methods for the presence of EE-GH1 in biological samples based on the use of such specific primers or probes.

The invention thus also relates to a kit for identifying elite event EE-GH1 in biological samples, the kit comprising at least one primer or probe which specifically recognizes the 5' or 3' flanking region of EE-GH1.

Preferably the kit of the invention comprises, in addition to a primer which specifically recognizes the 5' or 3' flanking region of EE-GH1, a second primer which specifically recognizes a sequence within the foreign DNA of EE-GH1, for use in a PCR identification protocol. Preferably, the kit of the invention comprises two (or more) specific primers, one of which recognizes a sequence within the 5' flanking region of EE-GH1, most preferably a sequence within the plant DNA region of SEQ ID NO: 1, and an other which recognizes a sequence within the foreign DNA. Especially preferably, the primer recognizing the plant DNA sequence within 5' flanking region comprises the nucleotide sequence of SEQ ID NO: 2. Particularly, the primer recognizing the plant DNA sequence within 5' flanking region comprises the nucleotide sequence of SEQ ID NO: 2 and the primer recognizing the foreign DNA comprises the nucleotide sequence of SEQ ID NO: 1 described herein.

According to a further embodiment, the invention relates to a kit for identifying elite event EE-GH1 in biological samples, which kit comprises at least one specific primer or probe having a sequence which corresponds (or is complementary to) a sequence that hybridizes under stringent conditions to a specific or characterizing region of EE-GH1.

Preferably the sequence of the probe corresponds to a specific region comprising part of the foreign DNA and part of the 5' or 3' flanking region (plant DNA) of EE-GH1 contiguous therewith. Most preferably the specific probe comprises (or is complementary to) a sequence that hybridizes under stringent conditions to a sequence of 15 to 30 nucleotides spanning the insertion site at the 5' or 3' region of the foreign DNA within SEQ ID NO: 1 or SEQ ID NO: 4. Especially preferably the specific probe comprises (or is complementary to) a sequence that hybridizes under stringent conditions to a sequence



of 9 nucleotides of plant DNA and a similar number of nucleotides of the foreign DNA contiguous with the insertion site, within the sequences of SEQ ID NO:1 or SEQ ID NO:4. According to a particular embodiment, such a primer comprises a sequence that is identical to a sequence of 9 nucleotides of plant DNA and a similar number of nucleotides of the foreign DNA contiguous with the insertion site, within the sequences of SEQ ID NO: 1 or SEQ ID NO: 4.

The methods and kits encompassed by the present invention can be used for different purposes such as, but not limited to the following: to identify EE-GH1 in plants, plant material or in products such as, but not limited to food or feed products (fresh or processed) comprising or derived from plant material; additionally or alternatively, the methods and kits of the present invention can be used to identify transgenic plant material for purposes of segregation between transgenic and non-transgenic material; additionally or alternatively, the methods and kits of the present invention can be used to determine the quality (i.e. percentage pure material) of plant material comprising EE-GH1.

The present invention further relates to a method for tracking plants comprising elite event EE-GH1 in their genome upon introduction into different cultivars.

It will be understood that particular embodiments of the invention are described by the dependent claims cited herein.

#### Brief description of the Figure

The following detailed description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying Figure, incorporated herein by reference, in which:

**Fig. 1. PCR analysis of other events and elite event EE-GH1 using the EE-GH1 PCR identification protocol.** Loading sequence of the gel: lane 1, molecular weight marker (100bp ladder), lane 2, DNA sample from a cotton plant comprising the transgenic event EE-GH1, lane 3, DNA samples from a cotton plant comprising another transgenic event, lane 4, DNA from wild-type cotton, lane 5, wild-type + 1 copy of the pGSV71-BamHI digest (positive control), lane 6, negative control (no template), lane 8, molecular weight marker (100bp ladder).

### Detailed Description of Preferred Embodiments

The term “gene” as used herein refers to any DNA sequence comprising several operably linked DNA fragments such as a promoter region, a 5’ untranslated region (the 5’UTR), a coding region (which may or may not code for a protein), and an untranslated 3’ region (3’UTR) comprising a polyadenylation site. Typically in plant cells, the 5’UTR, the coding region and the 3’UTR are transcribed into an RNA of which, in the case of a protein encoding gene, the coding region is translated into a protein. A gene may include additional DNA fragments such as, for example, introns. As used herein, a genetic locus is the position of a given gene in the genome of a plant.

The term “chimeric” when referring to a gene or DNA sequence is used to refer to the fact that the gene or DNA sequence comprises at least two functionally relevant DNA fragments (such as promoter, 5’UTR, coding region, 3’UTR, intron) that are not naturally associated with each other and/or originate, for example, from different sources. “Foreign” referring to a gene or DNA sequence with respect to a plant species is used to indicate that the gene or DNA sequence is not naturally found in that plant species, or is not naturally found in that genetic locus in that plant species. The term “foreign DNA” will be used herein to refer to a DNA sequence as it has incorporated into the genome of a plant as a result of transformation. The “transforming DNA” as used herein refers to a recombinant DNA molecule used for transformation. The transforming DNA usually comprises at least one “gene of interest” (e.g. a chimeric gene) which is capable of

conferring one or more specific characteristics to the transformed plant. The term “recombinant DNA molecule” is used to exemplify and thus can include an isolated nucleic acid molecule which can be DNA and which can be obtained through recombinant or other procedures.

5

As used herein the term “transgene” refers to a gene of interest (or the whole of the foreign DNA) as incorporated in the genome of a plant. A “transgenic plant” refers to a plant comprising at least one transgene in the genome of all of its cells.

- 10 The foreign DNA present in the plants of the present invention will preferably comprise a herbicide tolerance gene, more specifically a 35S-*bar* gene as the gene of interest.

A “herbicide tolerance” gene as used herein refers to a gene that renders the plant tolerant to a herbicide. An example of a herbicide tolerance gene is gene comprising a sequence  
15 encoding the enzyme phosphinothricin acetyl transferase, which detoxifies phosphinothricin, under the control of a constitutive promoter. More specifically, in the elite event of the present invention the herbicide tolerance gene comprises the coding sequence of the bialaphos resistance gene (*bar*) of *Streptomyces hygroscopicus* (Thompson *et al.* (1987) EMBO J 6: 2519-2523) under control of the 35S promoter from  
20 Cauliflower Mosaic Virus (Odell *et al.*, (1985), Nature 313: 810-812), also referred to as “35S-*bar*” herein. The expression of the 35S-*bar* gene confers tolerance to herbicidal compounds phosphinothricin or bialaphos or glufosinate, or more generally, glutamine synthase inhibitors, or salts or optical isomers thereof which will generally be referred to as “glufosinate tolerance” herein.

- 25 By hybridizing under “stringent conditions” is meant the conventional hybridizing conditions as described by Sambrook *et al.* (1989) (Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbour Laboratory Press, NY) which for instance can comprise the following steps: 1) immobilizing plant genomic DNA fragments on a filter, 2) prehybridizing the filter for 1 to 2 hours at 42°C in 50% formamide, 5 X SSPE,  
30 2 X Denhardt’s reagent and 0.1% SDS, or for 1 to 2 hours at 68°C in 6 X SSC, 2 X Denhardt’s reagent and 0.1% SDS, 3) adding the hybridization probe which has been

labeled, 4) incubating for 16 to 24 hours, 5) washing the filter for 20 min. at room temperature in 1X SSC, 0.1 %SDS, 6) washing the filter three times for 20 min. each at 68°C in 0.2 X SSC, 0.1 %SDS, and 7) exposing the filter for 24 to 48 hours to X-ray film at -70°C with an intensifying screen.

5

The incorporation of a recombinant DNA molecule in the plant genome typically results from transformation of a cell, tissue or callus (or from another genetic manipulation). The particular site of incorporation is either random or is at a predetermined location (if a process of targeted integration is used).

10

The DNA introduced into the plant genome as a result of transformation of a plant cell or tissue with a recombinant DNA or "transforming DNA" is hereinafter referred to as "foreign DNA" comprising one or more "transgenes". Thus, foreign DNA may comprise both recombinant DNA as well as newly introduced, rearranged DNA of the plant.

15

However, the term "plant DNA" in the context of the present invention will refer to DNA of the plant which is generally found in the same genetic locus in the corresponding wild-type plant.

20

The foreign DNA can be characterized by the location and the configuration at the site of incorporation of the recombinant DNA molecule in the plant genome. The site in the plant genome where a recombinant DNA has been inserted is also referred to as the "insertion site" or "target site". Insertion of the recombinant DNA into the plant genome can be associated with a deletion of plant DNA, referred to as "target site deletion". A "flanking region" or "flanking sequence" as used herein refers to a sequence of at least 20 bp, preferably at least 50 bp, and up to 5000 bp of the plant genome which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the foreign DNA. Transformation procedures leading to random integration of the foreign DNA will result in transformants with different flanking regions, which are characteristic and unique for each transformant. When the recombinant DNA present in a transgenic plant is introduced into a different plant through traditional crossing, its insertion site in the plant genome, or its flanking regions will generally not be changed (apart from occasional changes due to mutations or cross-

30

over and transposons). An “insertion region” as used herein refers to the region corresponding to the region of at least 40 bp, preferably at least 100 bp, and up to more than 10000 bp, encompassed by the sequence which comprises the upstream and/or the downstream flanking region of a foreign DNA in the (untransformed) plant genome (and including the insertion site and possible target site deletion). Taking into consideration minor differences due to mutations within a species, an insertion region will retain at least 85%, preferably 90%, more preferably 95%, and most preferably 100% sequence identity with the sequence comprising the upstream and downstream flanking regions of the foreign DNA in a given plant of that species.

Expression of a gene of interest refers to the fact that the gene confers on the plant one or more phenotypic traits (e.g. herbicide tolerance) that were intended to be conferred by the introduction of the recombinant DNA molecule – the transforming DNA - used during transformation (on the basis of the structure and function of part or all of the gene(s) of interest).

An “event” is defined as a (artificial) genetic locus that, as a result of genetic engineering, carries a foreign DNA comprising at least one copy of the gene(s) of interest (also referred to as a transformation event). An event is characterized phenotypically by the expression of the transgenes. At the genetic level, an event is part of the genetic makeup of a plant. At the molecular level, an event is characterized by the restriction map (e.g. as determined by Southern blotting) and/or by the upstream and/or downstream flanking sequences of the foreign DNA, and/or the molecular configuration of the foreign DNA comprising the transgenes. Usually when transforming a plant cell, tissue or callus with a transforming DNA, a multitude of events are generated, each of which is unique.

An “elite event”, as used herein, is an event which is selected from a group of events, obtained by transformation with the same transforming DNA or by back-crossing with plants obtained by such transformation, based on the phenotypic expression and stability of the transgenes and the absence of negative impact on the agronomic characteristics of the plant comprising it (i.e., selected transformation event). Thus the criteria for elite

event selection are one or more, preferably two or more, advantageously all of the following:

a) That the presence of the foreign DNA in the plant does not compromise other desired characteristics of the plant, such as those relating to agronomic performance or commercial value;

b) That the event is characterized by a well defined molecular configuration which is stably inherited and for which appropriate diagnostic tools for identity control can be developed;

c) That the gene(s) of interest show(s) an appropriate and stable spatial and temporal phenotypic expression in homozygous condition of the event, at a commercially acceptable level in a range of environmental conditions in which the plants carrying the event are likely to be exposed in normal agronomic use.

It is preferred that the foreign DNA is associated with a position in the plant genome that allows introgression into desired commercial genetic backgrounds.

The status of an event as an elite event is confirmed by introgression of the elite event in different relevant genetic backgrounds and observing compliance with one, two or all of the criteria e.g. a), b) and c) above.

An "elite event" thus refers to a genetic locus comprising a foreign DNA, which answers to the above-described criteria. A plant, plant material or progeny such as seeds can comprise one or more elite events in its genome. Thus, when referring to a plant, seed cell or tissue comprising elite event EE-GH1 in its genome, a plant, seed cell or tissue is intended which comprises the foreign DNA described herein (comprising the 35S-*bar* gene) integrated in its genome at the integration site described herein.

The tools developed to identify an elite event or the plant or plant material comprising an elite event, or products which comprise plant material comprising the elite event are based on the specific genomic characteristics of the elite event, such as, a specific restriction map of the genomic region comprising the foreign DNA, molecular markers or the sequence of the flanking region(s) of the foreign DNA.

Once one or both of the flanking regions of the foreign DNA have been sequenced, primers and probes can be developed which specifically recognize this (these) sequence(s) in the nucleic acid (DNA or RNA) of a sample by way of a molecular biological technique. For instance a PCR method can be developed to identify the elite event in biological samples (such as samples of plants, plant material or products comprising plant material). Such a PCR is based on at least two specific “primers” one recognizing a sequence within the 5’ or 3’ flanking region of the elite event and the other recognizing a sequence within the foreign DNA. The primers preferably have a sequence of between 15 and 35 nucleotides which under optimized PCR conditions “specifically recognize” a sequence within the 5’ or 3’ flanking region of the elite event and the foreign DNA of the elite event respectively, so that a specific fragment (“integration fragment”) is amplified from a nucleic acid sample comprising the elite event. This means that only the targeted integration fragment, and no other sequence (of that size) in the plant genome or foreign DNA, is amplified under optimized PCR conditions.

Preferably, the integration fragment has a length of between 50 and 500 nucleotides, most preferably of between 100 and 350 nucleotides. Preferably the specific primers have a sequence which is between 80 and 100% identical to a sequence within the 5’ or 3’ flanking region of the elite event and the foreign DNA of the elite event, respectively, provided the mismatches still allow specific identification of the elite event with these primers under optimized PCR conditions. The range of allowable mismatches however, can easily be determined experimentally and are known to a person skilled in the art.

As the sequence of the primers and their recognized sequence in the genome are unique for the elite event, amplification of the integration fragment will occur only in biological samples comprising (the nucleic acid of) the elite event. Preferably when performing a PCR to identify the presence of EE-GH1 in unknown samples, a control is included of a set of primers with which a fragment within a “housekeeping gene” of the plant species of the event can be amplified. Housekeeping genes are genes that are expressed in most cell types and which are concerned with basic metabolic activities common to all cells.

Preferably, the fragment amplified from the housekeeping gene is a fragment which is

larger than the amplified integration fragment. Depending on the samples to be analyzed, other controls can be included.

Standard PCR protocols are described in the art, such as in "PCR Applications Manual" (Roche Molecular Biochemicals, 2nd Edition, 1999). The optimal conditions for the PCR, including the sequence of the specific primers, is specified in a "PCR identification protocol" for each elite event. It is however understood that a number of parameters in the PCR identification protocol may need to be adjusted to specific laboratory conditions, and may be modified slightly to obtain similar results. For instance, use of a different method for preparation of DNA may require adjustment of, for instance, the amount of primers, polymerase and annealing conditions used. Similarly, the selection of other primers may dictate other optimal conditions for the PCR identification protocol. These adjustments will however be apparent to a person skilled in the art, and are furthermore detailed in current PCR application manuals such as the one cited above.

Alternatively, specific primers can be used to amplify an integration fragment that can be used as a "specific probe" for identifying EE-GH1 in biological samples. Contacting nucleic acid of a biological sample, with the probe, under conditions which allow hybridization of the probe with its corresponding fragment in the nucleic acid, results in the formation of a nucleic acid/probe hybrid. The formation of this hybrid can be detected (e.g. labeling of the nucleic acid or probe), whereby the formation of this hybrid indicates the presence of EE-GH1. Such identification methods based on hybridization with a specific probe (either on a solid phase carrier or in solution) have been described in the art. The specific probe is preferably a sequence which, under optimized conditions, hybridizes specifically to a region comprising part of the 5' or 3' flanking region of the elite event and also comprising part of the foreign DNA contiguous therewith (hereinafter also referred to as a "specific region" of the event). Depending on the method used, such a specific probe can comprise a sequence of between 15 and 30 bp or of between 50 and 500 bp, preferably of 100 to 350 bp which hybridizes under stringent conditions to the nucleotide sequence (or the complement of such sequence) of a specific region. Preferably, the specific probe will comprise a sequence of about 15 to about 100



contiguous nucleotides identical (or complementary) to a specific region of the elite event. Most preferably such a probe comprises at least 9 nucleotides identical to (or complementary to the 3' or 5' flanking region and a similar number of nucleotides in the foreign DNA contiguous therewith. Most preferably, such a probe comprises at least 9 nucleotides identical to (or complementary to the 3' or 5' flanking region and a similar number of nucleotides in the foreign DNA contiguous therewith of SEQ ID NO: 1 or SEQ ID NO: 4.

A "restriction map" as used herein refers to a set of Southern blot patterns obtained after cleaving plant genomic DNA (and/or the foreign DNA comprised therein) with a particular restriction enzyme, or set of restriction enzymes and hybridization with a probe sharing sequence similarity with the foreign DNA under standard stringency conditions. Standard stringency conditions as used herein refers to the conditions for hybridization described herein or to the conventional hybridizing conditions as described by Sambrook *et al.* (1989) (Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbour Laboratory Press, NY) which for instance can comprise the following steps: 1) immobilizing plant genomic DNA fragments on a filter, 2) prehybridizing the filter for 1 to 2 hours at 42°C in 50% formamide, 5 X SSPE, 2 X Denhardt's reagent and 0.1% SDS, or for 1 to 2 hours at 68°C in 6 X SSC, 2 X Denhardt's reagent and 0.1% SDS, 3) adding the hybridization probe which has been labeled, 4) incubating for 16 to 24 hours, 5) washing the filter for 20 min. at room temperature in 1X SSC, 0.1 %SDS, 6) washing the filter three times for 20 min. each at 68°C in 0.2 X SSC, 0.1 %SDS, and 7) exposing the filter for 24 to 48 hours to X-ray film at -70°C with an intensifying screen.

Due to the (endogenous) restriction sites present in a plant genome prior to incorporation of the foreign DNA, insertion of a foreign DNA will alter the specific restriction map of that genome. Thus, a particular transformant or progeny derived thereof can be identified by one or more specific restriction patterns.

Alternatively, plants or plant material comprising an elite event can be identified by testing according to a PCR identification protocol. This is a PCR which specifically

recognizes the elite event. Essentially, a set of PCR primers is developed which recognizes a) a sequence within the 3' or 5' flanking sequence of the elite event and b) a sequence within the foreign DNA, which primers amplify a fragment (integration fragment) preferably of between 100 and 300 nucleotides. Preferably, a control is included of a set of primers which amplifies a fragment within a housekeeping gene of the plant species (preferably a fragment which is larger than the amplified integration fragment). The optimal conditions for the PCR, including the sequence of the specific primers is specified in a PCR identification protocol.

Other methods for identifying plants, plant material or products comprising plant material comprising elite event EE-GH1 are also envisaged. These methods include all methods based on the detection of the foreign DNA sequence and flanking sequence(s) of the elite event with a specific probe. More particularly, chip-based technologies, such as those described by Hacia *et al.* 1996 (Nat Genet 14(4):441-447) and Shoemaker *et al.* 1996 (Nat Genet 14(4):450-456) are envisaged. These methods allow segregation of target molecules as high-density arrays by using fixed probe arrays or by tagging of the genes with oligonucleotides, after which they can be screened by hybridization. Alternatively, third wave detection methods, such as those described by Lyamichev *et al.* (1999, above) are envisaged.

Identification of the protein(s) encoded by the foreign DNA of the elite event can be done by classical protein detection methods described in the art, such as those based on chromatographic or electromagnetic properties of the protein or the detection by specific monoclonal antibodies (as described in "Guide to protein purification, Murray P. Deutscher editor).

A "kit" as used herein refers to a set of reagents for the purpose of performing the method of the invention, more particularly, the identification of the elite event EE-GH1 in biological samples. More particularly, a preferred embodiment of the kit of the invention comprises at least one or two specific primers, as described above. Optionally, the kit can further comprise any other reagent described herein in the PCR identification protocol.

Alternatively, according to another embodiment of this invention, the kit can comprise a specific probe, as described above, which specifically hybridizes with nucleic acid of biological samples to identify the presence of EE-GH1 therein. Optionally, the kit can further comprise any other reagent (such as but not limited to hybridizing buffer, label) for identification of EE-GH1 in biological samples, using the specific probe.

The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g., purity of seed lots), detection of the elite event in plant material or material comprising or derived from plant material, such as but not limited to food or feed products.

The present invention relates to the development of an elite event in cotton, EE-GH1, to the plants comprising this event, the progeny obtained from these plants and to the plant cells, or plant material derived from this event. Plants comprising elite event EE-GH1 were obtained through transformation with pGSV71 as described in example 1.

Cotton plants or plant material comprising EE-GH1 can be identified according to the PCR identification protocol described for EE-GH1 in Example 4 herein. Briefly, cotton genomic DNA is amplified by PCR using a primer which specifically recognizes a sequence within the 5' or 3' flanking sequence of EE-GH1, particularly the primer with the sequence of SEQ ID NO: 2, and a primer which recognizes a sequence in the foreign DNA, particularly the primer with the sequence of SEQ ID NO: 3. Endogenous cotton DNA primers are used as controls. If the plant material yields a fragment of between 250 and 290 bp, preferably of about 269 bp, the cotton plant is determined to harbor elite event EE-GH1.

Plants harboring EE-GH1 are characterized by their glufosinate tolerance, which in the context of the present invention includes that plants are tolerant to the herbicide Liberty<sup>TM</sup>. Tolerance to Liberty<sup>TM</sup> can be tested in different ways. The leaf paint method as described herein, is most useful when you wish to identify both resistant and sensitive plants, but do not want to kill the sensitive ones. Alternatively, tolerance can be tested by

Liberty<sup>TM</sup> spray application. Spray treatments should be made between the leaf stages V3 and V4 for best results. Tolerant plants are characterized by the fact that spraying of the plants with at least 200 grams active ingredient/hectare (g.a.i./ha), preferably 400 g.a.i./ha, and possibly up to 1600 g.a.i./ha (4X the normal field rate), does not kill the plants. A broadcast application should be applied at a rate of 28-34 oz Liberty<sup>TM</sup>. It is best to apply at a volume of 20 gallons of water per acre using a flat fan type nozzle while being careful not to direct spray applications directly into the whorl of the plants to avoid surfactant burn on the leaves. The herbicide effect should appear within 48 hours and be clearly visible within 5-7 days.

Plants harboring EE-GH1 can further be characterized by the presence in their cells of phosphinothricin acetyl transferase as determined by a PAT assay (De Block et al, 1987, supra).

Plants harboring EE-GH1 can, for example, be obtained from seeds deposited at the ATCC under accession number PTA-3343, which contain 50% kernels that are hemizygous for the elite event. Such plants can be further propagated to introduce the elite event of the invention into other cultivars of the same plant species. Selected seeds obtained from these plants contain the elite event stably incorporated into their genome.

The invention further relates to plants derived from the ATCC accession number PTA-334, comprising EE-GH1. The term "derived from" herein indicates that the plants are related, i.e. they are both progeny (direct or of two or more generations) of the same transformant by crossing.

Plants harboring EE-GH1 are also characterized by having agronomical characteristics that are comparable to commercially available varieties of cotton in the US, in the absence of weed pressure and use of Liberty<sup>TM</sup> for weed control. It has been observed that the presence of a foreign DNA in the insertion region of the cotton plant genome described herein, confers particularly interesting phenotypic and molecular characteristics to the plants comprising this event. More specifically, the presence of the foreign DNA in this particular region in the genome of these plants, results in plants which display a stable phenotypic expression of the gene of interest without significantly compromising

any aspect of desired agronomic performance of the plants. Thus, the insertion region, corresponding to a sequence comprising the plant DNA of SEQ ID NO: 1 and/or SEQ ID NO: 4, more particularly a sequence corresponding to SEQ ID NO: 5, most particularly the insertion site of EE-GH1 therein, is shown to be particularly suited for the introduction of a gene(s) of interest. More particularly, the insertion region of EE-GH1 (corresponding to a DNA sequence of at least 40 bp in the cotton genome within SEQ ID NO: 5), or a sequence of at least 40 bp which hybridizes under stringent conditions to the complement of the sequence of SEQ ID NO: 5, is particularly suited for the introduction of foreign DNA comprising a herbicide tolerance gene, ensuring expression of each of these genes in the plant without compromising agronomic performance.

A recombinant DNA molecule can be specifically inserted in an insertion region by targeted insertion methods. Such methods are well known to those skilled in the art and comprise, for example, homologous recombination using a recombinase such as, but not limited to the FLP recombinase from *Saccharomyces cerevisiae* (published PCTP application WO 99/25821), the CRE recombinase from *Escherichia coli* phage P1 (published PCT application WO 99/25840), the recombinase from pSR1 of *Saccharomyces rouxii* (Araki *et al.* 1985, J Mol Biol 182:191-203), the Gin/gix system of phage Mu (Maeser and Kahlmann, 1991, Mol Gen Genetics 230:170-176) or the lambda phage recombination system (such as described in US Patent 4,673,640).

As used herein, "sequence identity" with regard to nucleotide sequences (DNA or RNA), refers to the number of positions with identical nucleotides divided by the number of nucleotides in the shorter of the two sequences. The alignment of the two nucleotide sequences is performed by the Wilbur and Lipmann algorithm (Wilbur and Lipmann, 1983, Proc Nat Acad Sci USA 80:726) using a window-size of 20 nucleotides, a word length of 4 nucleotides, and a gap penalty of 4. Computer-assisted analysis and interpretation of sequence data, including sequence alignment as described above, can, e.g., be conveniently performed using the programs of the Wisconsin Package (from the Genetics Computer Group, Inc). Sequences are indicated as "essentially similar" when such sequences have a sequence identity of at least about 75%, particularly at least about

80%, more particularly at least about 85%, quite particularly about 90%, especially about 95%, more especially about 100%, quite especially are identical. It is clear that when RNA sequences are said to be essentially similar or have a certain degree of sequence identity with DNA sequences, thymine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence. "Complementary to" as used herein refers to the complementarity between the A and T (U), and G and C nucleotides in nucleotide sequences.

As used in herein, a "biological sample" is a sample of a plant, plant material or products comprising plant material. The term "plant" is intended to encompass cotton (such as but not limited to *Gossypium hirsutum*) plant tissues, at any stage of maturity, as well as any cells, tissues, or organs taken from or derived from any such plant, including without limitation, any seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts. "Plant material", as used herein refers to material which is obtained or derived from a plant. Products comprising plant material relate to food, feed or other products which are produced using plant material or can be contaminated by plant material. It is understood that, in the context of the present invention, such biological samples are preferably tested for the presence of nucleic acids specific for EE-GH1, implying the presence of nucleic acids in the samples. Thus the methods referred to herein for identifying elite event EE-GH1 in biological samples, preferably relate to the identification in biological samples of nucleic acids which comprise the elite event.

As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA sequence which is functionally or structurally defined, may comprise additional DNA sequences, etc.

The following examples describe the development and characteristics of cotton plants harboring the elite events EE-GH1 as well as the development of tools for the identification of elite event EE-GH1 in biological samples.

- 5 Unless otherwise stated, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbour Laboratory Press, NY and in Volumes 1 and 2 of Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA. Standard materials and methods for plant molecular work are
- 10 described in Plant Molecular Biology Labfax (1993) by R.D.D. Croy published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

In the description and examples, reference is made to the following sequences:

- |    |                |  |
|----|----------------|--|
|    | SEQ ID NO: 1:  | sequence comprising the 5' flanking region |
| 15 | SEQ ID NO: 2:  | primer GHI06                               |
|    | SEQ ID NO: 3:  | primer GHI05                               |
|    | SEQ ID NO: 4:  | sequence comprising the 3' flanking region |
|    | SEQ ID NO: 5:  | insertion region                           |
|    | SEQ ID NO: 6:  | plasmid pGSV71                             |
| 20 | SEQ ID NO: 7:  | plasmid pRVA44                             |
|    | SEQ ID NO: 8:  | primer MDB327                              |
|    | SEQ ID NO: 9:  | primer MLD015                              |
|    | SEQ ID NO: 10: | primer MLD016                              |
|    | SEQ ID NO: 11: | primer MDB612                              |
| 25 | SEQ ID NO: 12: | primer MDB053                              |
|    | SEQ ID NO: 13: | primer MDB356                              |
|    | SEQ ID NO: 14: | primer DPA017                              |
|    | SEQ ID NO: 15: | primer MLD019                              |
|    | SEQ ID NO: 16: | sequence comprising target site deletion   |
| 30 | SEQ ID NO: 17: | primer GHI01                               |
|    | SEQ ID NO: 18: | primer GHI02                               |

## EXAMPLES

Example 1: Transformation of cotton with a herbicide tolerance gene

5 **1.1. Construction of the chimeric DNA comprising the *bar* gene under the control of a constitutive promoter**

A plasmid pGSV71 was constructed following standard procedures. The sequence of the genetic elements of plasmid pGSV71 is given in table 1 (SEQ ID NO: 6):

10

Table 1. Nucleotide positions of the genetic elements in pGSV71

Nt positions	Abbreviation	Description and references
198-222	-	Right border Repeat from the TL-DNA from pTiB6S3 (Gielen <i>et al.</i> , 1984, EMBO J. 3:835-846)
223-249	-	Polylinker
250-1634	P35S3	Promoter of the 35S RNA of Cauliflower Mosaic Virus (Odell <i>et al.</i> , (1985), Nature 313: 810-812)
1635-2186	<i>bar</i>	Coding sequence encoding phosphinothricine-acetyl-transferase from <i>Streptomyces hygroscopicus</i> (Thompson <i>et al.</i> , (1987) EMBO J. 6: 2519-2523). The N-terminal two codons of the wild-type <i>bar</i> coding region have been substituted for codons ATG and GAC respectively.
2187-2205	-	Polylinker
2206-2465	3'nos	A 260 bp TaqI fragment from the 3'untranslated region of the nopaline-synthase gene originating from the T-DNA of pTiT37 (Depicker <i>et al.</i> , 1982, J. Mol. Appl. Genet. 1: 561-573 )
2466-2519	-	Polylinker
2520-2544	-	Left border Repeat from the TL-DNA from pTiB6S3 (Gielen <i>et al.</i> , 1984, EMBO J. 3:835-846)

**1.2. Transformation of *Gossypium hirsutum***

15

Cotton tissue from Coker312 plants was transformed with pGSV71 using *Agrobacterium* transformation (US 5,986,181) and regenerated to plants on appropriate media.



The small plantlets initiated on the selective regeneration media were transferred to new medium for germination (all medium is hormone-free). Plantlets were then transferred to the growth chambers or to the greenhouses.

- 5 Selection was done on phosphinothricin (PPT) at all stages except plantlet regeneration, which was done in the absence of PPT to accelerate growth. This resulted in a set of primary transformants (plants of generation T0).

10 Example 2: Development of events

**2.1. Development of lines carrying the event trait**

T0 shoots were transferred to greenhouse soil and plants were screened for glufosinate tolerance and for the presence of the PAT enzyme with a PAT ELISA (Steffens Biotechnische Analysen GmbH, Ebringen, Germany).

20 T1 to T3 plants were grown in the greenhouse and tested for Liberty<sup>TM</sup> tolerance at a 2x rate (by spraying 56 oz/ha). Positive plants were tested for expression of the bar gene using the Pat assay as described by Deblock et al. 1987 (EMBO J. 6:2513-2518).

Presence of the foreign DNA and copy number was checked by Southern blot analysis. Total genomic DNA was isolated from 1g of leaf tissue according to the CETAB method of Doyle *et al.* (1987, Phytochem. Bull. 19:11) and digested with EcoRI restriction enzyme.

Probes such as the following were used for Southern analysis :

“bar” probe: 474 bp KpnI-BglII digest of plasmid pDE110 (WO 92/09696)

“35S” probe: 892 bp NcoI-MunI digest of plasmid pRVA44 (SEQ ID NO: 7)

30 T2 Plants were also evaluated for general phenotypic characteristics compared to the non-transgenic isogenic lines. In later generations, the lines for which no negative penalties on phenotype or agronomic performance was observed for the presence of the transgene

either in hemizygous or in homozygous condition, as compared to wild-types were selected.

T4 material was grown in the field and tested under field conditions for Liberty™ tolerance according to different schedules.

In later generations, plants were compared to commercial varieties for yield, fiber quality and plant mapping data. Agronomic characteristics, such as plant height, height to node, boll retention, stand, vigor, fiber length, fiber strength and lint yield were evaluated.

It was determined that one event performed equally or better than the comparable checks and that for this event yield was dependent on background rather than on presence of the transgene.

## **2.2. Selection of an elite event**

This selection procedure, yielded one elite event which displayed optimal expression of the 35S-*bar* gene, i.e. tolerance to glufosinate ammonium, without penalty on agronomic performance and yield. This elite event was named EE-GH1.

## **2.3. Testing of EE-GH1 in cotton varieties with different genetic backgrounds and in different locations**

The selected event was introduced into different commercial genetic backgrounds, including FM989, FM 832, FM958, and FM966 and results of field trials of four different locations were compared. Plants were sprayed with 1600 g.a.i./ha, using different treatments (1x3-5 leaf stage, 4x, 3-5 leaf stage, 1x+1x, 3-5 leaf stage, 4x+4x, 3-5 leaf stage, 0 as control).

Seedling emergence and vigor rating for the elite event was very good.

No visible damage as a result of herbicide application was ever observed after application regardless of rate or stage of development at the time of application. There were no detrimental effects on morphology or growth habit of plants by herbicide application

- 5 Furthermore, the event had normal leaf, flower and boll morphology, excellent fertility, and showed no disease or abnormal insect susceptibility in multiple genetic backgrounds. During introgression into multiple genetic backgrounds no aberrant problems or abnormalities were observed over four generations.

## 10 **2.4. Genetic analysis of the locus**

The genetic stability of the insert for the EE-GH1 event was checked by molecular and phenotypic analysis in the progeny plants over several generations.

- 15 Southern blot analyses of plants of the T1, T2 and T3 generation were compared for the EE-GH1 event. The patterns obtained were found to be identical in the different generations. This proves that the molecular configuration of the foreign DNA in EE-GH1 was stable.

- 20 The EE-GH1 event displayed Mendelian segregation for the transgene as a single genetic locus in at least three subsequent generations indicating that the insert is stable.

### Example 3: Characterization of elite event EE-GH1

## 25 **3.1 In-depth molecular and genetic analysis of the locus**

- 30 Once the EE-GH1 event was identified as the event in which expression of the transgene as well as overall agronomic performance were optimal, the locus of the transgene was analyzed in detail on a molecular level. This included sequencing of the flanking regions of the transgene.

The sequence of the regions flanking the inserted transgene in the EE-GH1 event was determined using the TAIL-PCR protocol as described by Liu *et al.* (1995, Plant J. 8(3): 457-463).

5 a) Determination of the 5' flanking region

The primers used were:

		Sequence (5' → 3')	Position in pGSV71
Degenerate primer	MDB327	NTg.Agg.WTC.NWg.TSA.T (SEQ ID NO: 8)	-
Primary TAIL	MLD015	Tgg.TTC.CTA.gCg.TgA.gCC.AgT.g (SEQ ID NO: 9)	606→585
Second. TAIL	MLD016	AgC.TgC.TgC.TCT.TgC.CTC.TgT (SEQ ID NO: 10)	467→447
Tertiary TAIL	GHI05	ggA.CCg.TTA.TAC.ACA.ACg.Tag (SEQ ID NO: 3)	358→338

Whereby N=A,C, T or g; S=C or g; W=A or T

10

The fragment amplified using MDB327-GHI05 was ca. 1200 bp which was sequenced (5' flank: SEQ ID NO: 1). The sequence between bp 1 and bp 677 comprised plant DNA, while the sequence between bp 678 and bp 850 corresponded to pGSV71 DNA.

15 b) Determination of the 3' flanking region

The primers used were:

		Sequence (5' → 3')	Position in pGSV71
Degenerate primer	MDB612	NgT.gCT.SWg.ANA.WgA.T (SEQ ID NO: 11)	-
Primary TAIL	MDB053	CAT.gAC.gTg.ggT.TCC.Tgg.Cag.C (SEQ ID NO: 12)	2109-2130
Secondary TAIL	MDB356	AAT.CCT.gTT.gCC.ggT.CTT.gCg (SEQ ID NO: 13)	2252-2272
Tertiary TAIL	DPA017	gAT.TAg.AgT.CCC.gCA.ATT.ATA.C (SEQ ID NO: 14)	2362-2383

Whereby: N = A,C,T or g; S = C or g; W = A or T

The fragment amplified using MDB612-DPA017 was ca. 400 bp, the complete sequence of which was determined (SEQ ID NO: 4). The sequence between nucleotide 1 and 179 corresponds to T-DNA, while the sequence between nucleotide 180 and 426 corresponds to plant DNA.

#### c) Identification of the target site deletion

Using primers corresponding to sequences within the flanking regions of the transgene on the wildtype *Gossypium hirsutum* as a template, the insertion site of the transgene was identified.

The following primers were used:

	Sequence (5' → 3')	Position in 5'flank (SEQ ID NO: 1)	Position in 3'flank (SEQ ID NO: 4)
GHI06	TTg.CAC.CAT.CTA.gCT.CAC.TC (SEQ ID NO: 2)	815 → 795	-----

MLD019 CAA.gAT.gCg.AgC.AAC.TAT.gT (SEQ ID NO: 15) 285 → 266

This yielded a 200 bp fragment (SEQ ID NO: 16) in which bp 85 to 122 corresponds to a target site deletion.

5 Thus, the insertion region (SEQ ID NO: 5) as sequenced comprises:

1-677:	5' flanking region	bp 1 to 677 of SEQ ID NO: 1
678-714:	target site deletion	bp 85 to 122 of SEQ ID NO: 16
715-916:	3' flanking region	bp 180 to 426 of SEQ ID NO: 4

### 10 3.2. Genetic analysis of the locus

The genetic stability of the insert was checked by molecular and phenotypic analysis in the progeny plants over several generations.

15 Southern blot analyses on glufosinate tolerant plants of EE-GH1 cotton plants of the T<sub>0</sub>, T<sub>1</sub> and T<sub>2</sub> generation were compared and were found to be identical. This proves that the molecular configuration of the transgene in EE-GH1 containing plants was stable.

The EE-GH1 event displayed Mendelian segregation for the transgene as a single genetic locus in at least three subsequent generations indicating that the insert is stable.

20

On the basis of the above results EE-GH1 was identified as an elite event.

#### Example 4: Development of diagnostic tools for identity control

25

A EE-GH1 Elite event PCR Identification protocol was developed to identify the presence of EE-GH1 in plants, plant material or biological samples.

**EE-GH1 Elite event Polymerase Chain Reaction identification protocol**

A test run, with all appropriate controls, has to be performed before attempting to screen unknowns. The presented protocol might require optimization for components that may differ between labs (template DNA preparation, Taq DNA polymerase, quality of the primers, dNTP's, thermocycler, etc.).

Amplification of the endogenous sequence plays a key role in the protocol. One has to attain PCR and thermocycling conditions that amplify equimolar quantities of both the endogenous and transgenic sequence in a known transgenic genomic DNA template. Whenever the targeted endogenous fragment is not amplified or whenever the targeted sequences are not amplified with the same ethidium bromide staining intensities, as judged by agarose gel electrophoresis, optimization of the PCR conditions may be required.

**Template DNA**

Template DNA is prepared according to the CTAB method described by Doyle and Doyle (1987, Phytochem. Bull. 19: 11). When using DNA prepared with other methods, a test run utilizing different amounts of template should be done. Usually 50 ng of genomic template DNA yields the best results.

**Assigned positive and negative controls**

The following positive and negative controls should be included in a PCR run:

- Master Mix control (DNA negative control). This is a PCR in which no DNA is added to the reaction. When the expected result, no PCR products, is observed this indicates that the PCR cocktail was not contaminated with target DNA.

- A DNA positive control (genomic DNA sample known to contain the transgenic sequences). Successful amplification of this positive control demonstrates that the PCR was run under conditions which allow for the amplification of target sequences.

- 5 - A wildtype DNA control. This is a PCR in which the template DNA provided is genomic DNA prepared from a non-transgenic plant. When the expected result, no amplification of the transgene PCR product but amplification of the endogenous PCR product, is observed this indicates that there is no detectable transgene background amplification in a genomic DNA sample.

10

### Primers

The following primers, which specifically recognize the transgene and a flanking sequence of EE-GH1 are used:

15

Primer	Sequence (5' → 3')	Position in SEQ ID NO: 1	Target
GHI05	ggA.CCg.TTA.TAC.ACA.ACg.Tag (SEQ ID NO: 3)	758→ 738	pGSV71 sequence
GHI06	TTg.CAC.CAT.CTA.gCT.CAC.TC (SEQ ID NO: 2)	815→ 795	Plant DNA Sequence

Primers targeting an endogenous sequence are always included in the PCR cocktail. These primers serve as an internal control in unknown samples and in the DNA positive control. A positive result with the endogenous primer-pair demonstrates that there is ample DNA of adequate quality in the genomic DNA preparation for a PCR product to be generated. The endogenous primers used are:

20

GHI01: 5'-AAC.CTA.ggC.TgC.TgA.Agg.AgC-3' (SEQ ID NO: 17)  
(Alcohol dehydrogenase gene Acc. NO: AF036569, 1070→1090)

25



GHI02: 5'-CAA.CTC.CTC.CAg.TCA.TCT.CCg-3' (SEQ ID NO: 18)  
 (Alcohol dehydrogenase gene Acc. NO: AF036569, 1515→1495)

### Amplified fragments

5

The expected amplified fragments in the PCR reaction are:

For primer pair GHI01-GHI02: 445bp (endogenous control)

For primer pair GHI05-GHI06: 269bp (EE-GH1 Elite Event)

10

### PCR conditions

The PCR mix for 50µl reactions contains:

5 µl template DNA  
 5 µl 10x Amplification Buffer (supplied with Taq polymerase)  
 1 µl 10 mM dNTP's  
 0.5 µl GHI01 (10pmoles/µl)  
 0.5 µl GHI02 (10pmoles/µl)  
 1 µl GHI05 (10pmoles/µl)  
 1 µl GHI06 (10pmoles/µl)  
 0.2 µl Taq DNA polymerase (5 units/µl)  
 water up to 50 µl

The thermocycling profile to be followed for optimal results is the following:

4 min. at 95°C  
 Followed by: 1 min. at 95°C  
 1 min. at 57°C  
 2 min. at 72°C  
 For 5 cycles  
 Followed by: 30 sec. at 92°C  
 30 sec. at 57°C

1 min. at 72°C

For 25 cycles

Followed by: 5 minutes at 72°C

5 **Agarose gel analysis .**

Between 10 and 20µl of the PCR samples should be applied on a 1.5% agarose gel (Tris-borate buffer) with an appropriate molecular weight marker (e.g. 100bp ladder PHARMACIA).

10

**Validation of the results**

Data from transgenic plant DNA samples within a single PCR run and a single PCR cocktail should not be acceptable unless 1) the DNA positive control shows the expected  
15 PCR products (transgenic and endogenous fragments), 2) the DNA negative control is negative for PCR amplification (no fragments) and 3) the wild-type DNA control shows the expected result (endogenous fragment amplification).

Lanes showing visible amounts of the transgenic and endogenous PCR products of the  
20 expected sizes, indicate that the corresponding plant from which the genomic template DNA was prepared, has inherited the EE-GH1 elite event. Lanes not showing visible amounts of the transgenic PCR product and showing visible amounts of the endogenous PCR product, indicate that the corresponding plant from which the genomic template DNA was prepared, does not comprise the elite event. Lanes not showing visible amounts  
25 of the endogenous and transgenic PCR products, indicate that the quality and/or quantity of the genomic DNA didn't allow for a PCR product to be generated. These plants cannot be scored. The genomic DNA preparation should be repeated and a new PCR run, with the appropriate controls, has to be performed.

**Use of discriminating PCR protocol to identify EE-GH1**

Cotton leaf material from plants comprising different transgenic events (samples 1 to 4) was tested according to the above-described protocol. Samples from cotton wild-type were taken as negative controls.

The results of the PCR analysis are illustrated in Figure 1. Sample 1 is recognized as comprising elite event EE-GH1. All other tested lines do not comprise this elite event.

**Example 5: Introgression of EE-GH1 into preferred cultivars**

Elite event EE-GH1 is introduced by repeated back-crossing into the following commercial cotton cultivars: FM5013, FM5015, FM5017, FM989, FM832, FM966 and FM958.

It is observed that the introgression of the elite event into these cultivars does not significantly influence any of the desirable phenotypic or agronomic characteristics of these cultivars (no linkage drag) while expression of the transgene, as determined by glufosinate tolerance, meets commercially acceptable levels. This confirms the status of event EE-GH1 as an elite event.

As used in the claims below, unless otherwise clearly indicated, the term “plant” is intended to encompass plant tissues, at any stage of maturity, as well as any cells, tissues, or organs taken from or derived from any such plant, including without limitation, any seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts.

Reference seed comprising elite event EE-GH1 was deposited as EE-GH1 at the ATCC (10801 University Blvd., Manassas, VA 20110-2209) on April 26, 2001, under ATCC accession number PTA-3343.

As used in the claims below, unless otherwise clearly indicated, the term “plant” is intended to encompass plant tissues, at any stage of maturity, as well as any cells, tissues, or organs taken from or derived from any such plant, including without limitation, any  
5 seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts.

The above description of the invention is intended to be illustrative and not limiting. Various changes or modifications in the embodiments described may occur to those skilled  
10 in the art. These can be made without departing from the spirit or scope of the invention.

## WHAT IS CLAIMED IS:

1. A transgenic cotton plant or seed, cells or tissues thereof comprising event EE-GH1 in its genome.
2. The transgenic cotton plant, or seed, cells or tissues of claim 1, the genomic DNA of which, when analyzed using the Elite event identification protocol for EE-GH1 with two primers comprising the nucleotide sequence of SEQ ID NO: 2 and SEQ ID NO: 3 respectively, yields a DNA fragment of between 250 and 290 bp.
3. The cotton plant, or seed, cells or tissues thereof, according to claim 2, wherein said DNA fragment is a fragment of about 269 bp.
4. A cotton plant, or seed, cells or tissues thereof, obtained by propagation of and/or breeding with a cotton plant grown from the seed deposited at the ATCC under accession number PTA-3343.
5. A cotton plant, seed, cells or tissues thereof which is the progeny of the seed deposited at the ATCC under accession number PTA-3343.
6. A method for identifying elite event EE-GH1 in biological samples, which method comprises detecting an EE-GH1 specific region with a primer or probe which specifically recognizes the 5' flanking region of SEQ ID NO: 3 or the 3' flanking region of SEQ ID NO: 4 of EE-GH1.
7. The method of claim 6, said method comprising amplifying a DNA fragment of between 100 and 350 bp from a nucleic acid present in said biological samples using a polymerase chain reaction with at least two primers, one of which recognizes the 5' or 3' flanking region of EE-GH1 and the other which recognizes a sequence within the foreign DNA of EE-GH1.

8. The method of claim 7, wherein said one primer recognizes a sequence within the 5' flanking region of SEQ ID NO: 3 and said other primer recognizes a sequence within the foreign DNA of EE-GH1.

9. The method of claim 8, wherein said primer recognizing a sequence within the 5' flanking region of EE-GH1 comprises the sequence of SEQ ID NO: 2.

10. The method of any one of claims 6 to 9, wherein said primer recognizing a sequence within the foreign DNA comprises the sequence of SEQ ID NO: 1.

11. A method for identifying EE-GH1 in a biological sample, which method comprises detecting an EE-GH1 specific region with a specific primer or probe which hybridizes under stringent conditions to a sequence within the 5' of SEQ ID NO: 3 or within the 3' flanking sequence of SEQ ID NO: 4 of EE-GH1.

12. A method for identifying a transgenic plant, or cells or tissues thereof, comprising the elite event EE-GH1, which method comprises establishing that genomic DNA can be used, according to a PCR identification protocol, to amplify a DNA fragment of between 250 and 290 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO: 1 and SEQ ID NO: 2, respectively.

13. A kit for identifying elite event EE-GH1 in biological samples, said kit comprising at least one PCR primer or probe, which recognizes a sequence within the 5' flanking region of SEQ ID NO: 3 or the 3' flanking region of SEQ ID NO: 4 of EE-GH1.

14. The kit of claim 13, wherein said at least one PCR primer recognizes a sequence within the plant DNA in SEQ ID NO: 3.

15. The kit of claim 14, wherein said primer recognizing a sequence within the plant DNA in SEQ ID NO: 3 comprises the sequence of SEQ ID NO: 2.

16. The kit of Claims 13 to 15, which further comprises at least a second PCR primer or probe which recognizes a sequence within the foreign DNA of EE-GH1.

17. The kit of claim 16, wherein said primer recognizing a sequence within the foreign DNA of EE-GH1 comprises the sequence of SEQ ID NO: 1.

18. A method for confirming seed purity, which method comprises detecting an EE-GH1 specific DNA sequence with a specific primer or probe which specifically recognizes a sequence within the 5' flanking region of SEQ ID NO: 3 or the 3' flanking region of SEQ ID NO: 4 of EE-GH1, in seed samples.

19. A method for screening seeds for the presence of EE-GH1, which method comprises detecting an EE-GH1 specific DNA sequence with a specific primer or probe which specifically recognizes a sequence within the 5' flanking region of SEQ ID NO: 3 or the 3' flanking region of SEQ ID NO: 4 of EE-GH1, in samples of seed lots.

20. A seed deposited at the ATCC under accession number PTA-3343.

21. A cotton seed comprising elite event EE-GH1, reference seed comprising said event having been deposited at the ATCC under accession number PTA-3343.

22. A cotton plant, cell or tissue or plant material thereof comprising elite event EE-GH1, derived from the seed of claim 21.

23. Transgenic cotton plants, seeds, cells or tissues, the genomic DNA of which comprises a transgene integrated into the chromosomal DNA in a region which

comprises a sequence of at least 40 bp which hybridizes under stringent conditions with a sequence which is complementary to the sequence of SEQ ID NO: 5.

24. A process for producing a transgenic cotton plant or cell or tissue of a cotton plant, said process comprising introducing a recombinant DNA molecule into a region of cotton chromosomal DNA corresponding to a sequence of at least 40 bp that hybridizes under stringent conditions with a sequence that is complementary to the sequence of SEQ ID NO: 5, and, optionally, regenerating a cotton plant from the transformed cotton cell or tissue.

25. The process of claim 24, wherein said recombinant DNA molecule comprises an herbicide resistance gene.

26. The plant or cell or tissue of a cotton plant obtained by the process of claims 24 or 25.

27. A transgenic cotton plant or seed, cells or tissues thereof comprising  
(i) event EE-GH1 in its genome ;or  
(ii) event EE-GH1 with the proviso that the *bar* gene used in the event is substituted with a nucleic acid sequence that hybridizes to the complement of the *bar* gene under stringent conditions.

28. A DNA molecule which can be amplified from a nucleic acid sample of cotton, using a set of primers the first of which comprises a sequence of 15 to 30 nucleotides complementary to a sequence within SEQ ID NO:1 or SEQ ID NO:4, the second of which comprises a sequence complementary to the foreign DNA present in said nucleic acid sample of corn.

29. The DNA molecule of claim 28, which can be amplified from a nucleic acid sample of cotton, using a set of primers, the first of which comprising SEQ ID NO:2, the second of which comprising SEQ ID NO:3.



30. A detection kit comprising the DNA of claim 28 or 29.
31. A DNA molecule isolated from corn tissue, comprising the sequence of SEQ ID  
5 NO:1 or SEQ ID NO:4
32. A DNA molecule comprising a sequence which is identical to or complementary to  
the sequence of 9 nucleotides of plant DNA and foreign DNA at each side  
contiguous with the insertion site contained in SEQ ID NO: 1 or SEQ ID NO: 4.

10




Figure 1

## SEQUENCE LISTING

<110> Aventis CropScience N.V.

<120> Herbicide tolerant cotton plants and methods for producing and identifying same

<130> EE-GH1

<150> US 09/921,922

<151> 2001-08-06

<160> 18

<170> PatentIn version 3.1

<210> 1

<211> 850

<212> DNA

<213> Artificial: sequence comprising 5' flanking region

<400> 1  
aaaggggatg agattgaatg ttaccttattc aacaaaagga gttgtagctc atggaacaac 60  
aatagtcttt tccacggaaa cctagatgat gtttctccaa tgcttgataa atctttaaca 120  
ttgtcatcat aagttgcaac ctcatgtttc acacaagcat caatcaaattg ttgatcttca 180  
ttactaaaat gtgcttgatc cttccttaca caaatctacc tatgttggtg tattttgttc 240  
tattcatcat tctaacaagt ttgcaattg agttgaactt cttccaatct cgtatcagcc 300  
tataatagtg ggggtctaata tgtccatttt tcccacaata atgacatata atctttctaa 360  
agctttttatt ctctgcctta tgatgaaaag aacccaaatc tttaacttta acaaaaataa 420  
gatgagcgat aggttcttca cctttattga tgtaaccaag tcctctatgg catggttcaa 480  
ttctcattga agccaaaatt tcatgaaact tctcacattg gcctctaaac ttcttcaaga 540  
tagcctttgc accatctagc tcactcttgg ttgttttcaa aacatcatcc gtttcttgga 600  
ccacaatttt gagcttttca ttttctattt tgaggataat agttttattcc ctcaaggaac 660  
tattcaactg agcttaacag tactcggccg tcgaccgagg taccoggaat tccaatccca 720  
caaaaatctg agcttaacag cacagttgct cctctcagag cagaatcggg tattcaacac 780  
cctcatatca actactacgt tgtgtataac ggtccacatg ccggtatata cgatgactgg 840  
ggttgtaaaa 850

<210> 2

<211> 20

<212> DNA

<213> Artificial: primer GHI06

<400> 2

ttgcaccatc tagctcactc

20

<210> 3  
 <211> 21  
 <212> DNA  
 <213> Artificial: primer GHI05

<400> 3  
 ggaccgttat acacaacgta g 21

<210> 4  
 <211> 426  
 <212> DNA  
 <213> Artificial: sequence comprising 3'flanking region

<400> 4  
 gattagagtc ccgcaattat acatttaata cgcgatagaa aacaaaatat agcgcgcaaa 60  
 ctaggataaa ttatcgcgcg cgggtgtcatc tatgttacta gatcggggaag atcctctaga 120  
 gtcgacctgc aggcattgcaa gcttagatcc atggagccat ttacaattga atatatcctc 180  
 caaatattta aaaagaatat caccattatc cgaatcttct ttaaaatctg ttagaacacg 240  
 gtttggaata gtggtagtaa aagtaacata gttgctcgca tcttgatcta cattaaactt 300  
 tcttcatcac tccaagtgat tgtaaatgac ttctatttct tcttagtatt agcacattct 360  
 aattttaagt gaaacaatcc cttacattca taacattgaa tctccttcta tcatctcaca 420  
 gcacga 426

<210> 5  
 <211> 961  
 <212> DNA  
 <213> Gossypium hirsutum: insertion region

<400> 5  
 aaaggggatg agattgaatg ttaccttatt aacaaaagga gttgtagctc atggaacaac 60  
 aatagtcttt tccacggaaa cctagatgat gtttctccaa tgcttgataa atctttaaca 120  
 ttgtcatcat aagttgcaac ctcatgtttc acacaagcat caatcaaagtg ttgatcttca 180  
 ttactaaaat gtgcttgatc cttccttaca caaatctacc tatgttggtgg tattttgttc 240  
 tattcatcat tctaacaagt tttgcaattg agttgaactt cttccaatct cgtatcagcc 300  
 tataatagtg gggcttaata tgtccatttt tcccacaata atgacatata atctttctaa 360  
 agctttttatt ctctgcctta tgatgaaaag aacccaaatc tttaacttta acaaaaaataa 420  
 gatgagcgat aggttcttca cttttattga tgtaaccaag tctctatagg catgggtcaa 480  
 ttctcattga agccaaaatt tcatgaaact tctcacattg gcctctaaac ttcttcaaga 540  
 tagcctttgc accatctagc tcaactcttg ttgttttcaa aacatcatcc gtttcttgga 600  
 ccacaatttt gagcttttca ttttctattt tgaggataat agtttattcc ctcaaggaac 660

tattcaactg agcttaaate tcaattttttt ttaacatatg actataagta tcctccaaat	720
atttaaaaag aatatcacca ttatccgaat cttcttttaa atctgttaga acacgggttg	780
gaatagtggg agtaaaagta acatagttgc tcgcatcttg atctacatta aactttcttc	840
atcactccaa gtgattgtaa atgacttcta tttcttctta gtattagcac attctaattt	900
taagtgaaac aatcccttac attcataaca ttgaatatcc ttctatcatc tcacagcacg	960
a	961

&lt;210&gt; 6

&lt;211&gt; 9555

&lt;212&gt; DNA

&lt;213&gt; Artificial: plasmid pGSV71

&lt;400&gt; 6

agattcgaag ctcggtcccg tgggtgttct gtcgtctcgt tgtacaacga aatccattcc	60
cattccgcgc tcaagatggc ttccccctcg cagttcatca gggctaaatc aatctagccg	120
acttgtcccg tgaaatgggc tgcactccaa cagaaacaat caaacaacaa tacacagcga	180
cttattcaca cgcgacaaat tacaacggta tatatcctgc cagtactcgg ccgtcgaccg	240
cggatcccg aattccaatc ccacaaaaat ctgagcttaa cagcacagtt gtcctctca	300
gagcagaatc gggatttcaa caccctcata tcaactacta cgttgtgtat aacgggtccac	360
atgccgggat atacgatgac tgggggttga caaaggcggc aacaaacggc gttcccggag	420
ttgcacacaa gaaatttgcc actattacag aggcaagagc agcagctgac gcgtacacaa	480
caagtcagca aacagacagg ttgaacttca tccccaaagg agaagctcaa ctcaagccca	540
agagctttgc taaggcccta acaagcccac caaagcaaaa agcccactgg ctcacgctag	600
gaacccaaaag gcccagcagt gatccagccc caaaagagat ctcctttgcc ccggagatta	660
caatggacga tttcctctat ctttacgac taggaaggaa gttcgaagggt gaagggtgacg	720
acactatggt caccactgat aatgagaagg tttagcctct caatttcaga aagaatgctg	780
acccacagat ggtagagag gcctacgcag cagggtctcat caagacgac taccgagta	840
acaatctcca ggagatcaaa taccttccca agaagggtta agatgcagtc aaaagattca	900
ggactaattg catcaagaac acagagaaag acatatttct caagatcaga agtactattc	960
cagtatggac gattcaaggc ttgcttcata aaccaaggca agtaatagag attggagtct	1020
ctaaaaaggt agttcctact gaatctaagg ccatgcatgg agtctaagat tcaaactcgag	1080
gatctaacag aactcgccgt gaagactggc gaacagttca tacagagtct ttacgactc	1140
aatgacaaga agaaaatctt cgtcaacatg gtggagcacg acactctggg ctactccaaa	1200
aatgtcaaag atacagtctc agaagaccaa agggctattg agacttttca acaaaggata	1260

atttcgggaa acctcctcgg attccattgc ccagctatct gtcacttcat cgaaaggaca	1320
gtagaaaagg aaggtggctc ctacaaatgc catcattgcg ataaaggaaa ggctatcatt	1380
caagatgcct ctgccgacag tggccccaaa gatggacccc caccacgag gagcatcgtg	1440
gaaaaagaag acgttccaac cacgtcttca aagcaagtgg attgatgtga catctccact	1500
gacgtaaggg atgacgcaca atcccactat ccttcgcaag acccttcctc tatataagga	1560
agttcatttc atttggagag gacacgctga aatcaccagt ctctctctat aaatctatct	1620
ctctctctat aaccatggac ccagaacgac gcccggccga catccgccgt gccaccgagg	1680
cggacatgcc ggcggtctgc accatcgtca accactacat cgagacaagc acggtcaact	1740
tccgtaccga gccgcaggaa ccgcaggagt ggacggacga cctcgtccgt ctgcgggagc	1800
gctatccctg gctcgtcgcc gaggtggacg gcgaggtcgc cggcatcgcc tacgcgggcc	1860
cctggaaggc acgcaacgcc tacgactgga cggccgagtc gaccgtgtac gtctcccccc	1920
gccaccagcg gacgggactg ggctccacgc tctacacca cctgctgaag tccctggagg	1980
cacagggctt caagagcgtg gtcgctgtca tcgggctgcc caacgaccg agcgtgcgca	2040
tgcacgaggc gctcggatat gcccccgcg gcatgctgcg ggcggccggc ttcaagcacg	2100
ggaactggca tgacgtgggt ttctggcagc tggacttcag cctgccggta ccgccccgtc	2160
cggtcctgcc cgtcaccgag atctgagatc acgcgttcta ggatccgaag cagatcgttc	2220
aaacatttgg caataaagtt tcttaagatt gaatcctggt gccggtcttg cgatgattat	2280
catataattt ctggtgaatt acgttaagca tgtaataatt aacatgtaat gcatgacggt	2340
atttatgaga tgggttttta tgattagagt cccgcaatta tacatttaat acgcgataga	2400
aaacaaaata tagcgcgcaa actaggataa attatcgcg cgggtgtcat ctatgttact	2460
agatcgggaa gatcctctag agtcgaactg caggcatgca agcttagatc catggagcca	2520
tttacaattg aatatatcct gccgcgcgtg ccgctttgca cccggtggag cttgcatggt	2580
ggtttctacg cagaactgag ccggttaggc agataatttc cattgagaac tgagccatgt	2640
gcaccttccc cccaacacgg tgagcgacgg ggcaacggag tgatccacat gggactttta	2700
aacatcatcc gtcggatggc gttgcgagag aagcagtcga tccgtgagat cagccgacgc	2760
accgggcagg cgcgcaacac gatcgcaaag tatttgaacg caggtacaat cgagccgacg	2820
ttcacggtag cggaacgacc aagcaagcta gcttagtaaa gccctcgcta gattttaatg	2880
cggatgttgc gattacttcg ccaactattg cgataacaag aaaaagccag cctttcatga	2940
tatatctccc aatttgtgta gggcttatta tgcacgctta aaaataataa aagcagactt	3000
gacctgatag tttggctgtg agcaattatg tgcttagtgc atctaacgt tgagttaagc	3060
cgcgccgcga agcggcgctc gcttgaacga attgttagac attatttgcc gactaccttg	3120

gtgatctcgc ctttcacgta gtggacaaat tcttccaact gatctgcgcg cgaggccaag 3180  
 cgatcttctt cttgtccaag ataagcctgt ctagcttcaa gtatgacggg ctgatactgg 3240  
 gccggcaggc gctccattgc ccagtcggca gcgacatcct tcggcgcgat tttgccggtt 3300  
 actgcgctgt accaaatgcg ggacaacgta agcactacat ttcgctcatc gccagcccag 3360  
 tcggggcgcg agttccatag cgttaagggtt tcatttagcg cctcaaatag atcctgttca 3420  
 ggaaccggat caaagagttc ctccgccgct ggacctacca aggcaacgct atgtttctct 3480  
 gcttttgtca gcaagatagc cagatcaatg tcgatcgtgg ctggctcgaa gataacctgca 3540  
 agaatgtcat tgcgctgcca ttctccaaat tgcagttcgc gcttagctgg ataacgccac 3600  
 ggaatgatgt cgtcgtgcac aacaatggtg acttctacag cgcggagaat ctgcctctct 3660  
 ccaggggaag ccgaagtctc caaaaggctg ttgatcaaag ctgcgccgct tgtttcatca 3720  
 agccttacgg tcaccgtaac cagcaaatac atatcactgt gtggcttcag gccgccatcc 3780  
 actgcggagc cgtacaaatg tacggccagc aacgtcgggt cgagatggcg ctcgatgacg 3840  
 ccaactacct ctgatagttg agtcgatact tcggcgatca ccgcttccct catgatgttt 3900  
 aactttgttt tagggcgact gccctgctgc gtaacatcgt tgctgctcca taacatcaaa 3960  
 catcgaccca cggcgtaacg cgcttgctgc ttggatgcc gaggcataga ctgtacccca 4020  
 aaaaaacagt cataacaagc catgaaaacc gccactgcgc cgttaccacc gctgcgttcg 4080  
 gtcaagggtt tggaccagtt gcgtgagcgc atacgtact tgcattacag cttacgaacc 4140  
 gaacaggctt atgtccactg ggttcgtgcc ttcacccggt tccacgggtg gcgtcaccgc 4200  
 gcaaccttgg gcagcagcga agtcgaggca tttctgtcct ggctggcgaa cgagcgcaag 4260  
 gtttcggtct ccacgcatcg tcaggcattg gcggccttgc tgttcttcta cggcaagtgc 4320  
 tgtgcacgga tctgccctgg cttcaggaga tcggaagacc tcggccgtcc gggcgcttgc 4380  
 cgggtggtgct gaccccggtg gaagtgggtc gcatcctcgg ttttctggaa ggcgagcatc 4440  
 gtttgttcgc ccagcttctg tatggaacgg gcatgcggat cagtgagggt ttgcaactgc 4500  
 ggggtcaagga tctggatttc gatcacggca cgatcatcgt gcgggagggc aagggtcca 4560  
 aggatcgggc cttgatgtta cccgagagct tggcaccag cctgcgcgag cagggatcga 4620  
 tccaacccct ccgctgctat agtgcagtcg gcttctgacg ttcagtgcag ccgtcttctg 4680  
 aaaacgacat gtcgcacaag tcctaagtta cgcgacaggc tgccgccctg cccttttctt 4740  
 ggcgttttct tgtcgcgtgt tttagtcgca taaagtagaa tacttgcgac tagaaccgga 4800  
 gacattacgc catgaacaag agcgcgcgcg ctggcctgct gggctatgcc cgcgtcagca 4860  
 ccgacgacca ggacttgacc aaccaacggg ccgaactgca cgcggccggc tgcaccaagc 4920

tgttttccga gaagatcacc ggcaccaggc ggcaccgccc ggagctggcc aggatgcttg	4980
accacctacg ccctggcgac gttgtgacag tgaccaggct agaccgcctg gcccgcagca	5040
cccgcgacct actggacatt gccgagcgca tccaggaggc cggcgcgggc ctgcgtagcc	5100
tggcagagcc gtggggccgac accaccacgc cggccggccg catggtgttg accgtgttcg	5160
ccggcattgc cgagttcgag cgttccttaa tcatcgaccg caccgcgagc gggcgcgagg	5220
ccgccaaagg ccgaggcgtg aagtttggcc cccgccctac cctcaccg gcacagatcg	5280
cgcacgcccc cgagctgacg gaccaggaag gccgcaccgt gaaagaggcg gctgcaactgc	5340
ttggcggtgca tcgctcgacc ctgtaccg cgacttgagcg cagcgaggaa gtgacgcca	5400
ccgaggccag gcggcgcggt gccttcctg aggacgcatt gaccgaggcc gacgccctgg	5460
cggccgcccga gaatgaacgc caagaggaac aagcatgaaa ccgcaccagg acggccagga	5520
cgaaccgttt ttcattaccg aagagatcga ggcggagatg atcgcgccg ggtacgtgtt	5580
cgagccgccc gcgcacgtct caaccgtgcg gctgcatgaa atcctggccg gtttgtctga	5640
tgccaagctg gcggcctggc cggccagctt ggccgctgaa gaaaccgagc gccgccgtct	5700
aaaaaggtga tgtgtatttg agtaaaacag cttgcgtcat gcggtcgctg cgtatatgat	5760
gcgatgagta aataaaciaa tacgcaaggg gaacgcacga aggttatcgc tgtacttaac	5820
cagaaaggcg ggtcaggcaa gacgaccatc gcaaccatc tagcccgcg cctgcaactc	5880
gccggggccg atgttctgtt agtcgattcc gatccccagg gcagtggccg cgattgggcg	5940
gccgtgcggg aagatcaacc gctaaccgtt gtcggcatcg accgcccgc gattgaccgc	6000
gacgtgaagg ccatcggccg gcgcgacttc gtagtgatcg acggagcgcc ccaggcgcg	6060
gacttggtcg tgtcccgat caaggcagcc gacttcgtgc tgattccggt gcagccaagc	6120
ccttacgaca tatgggccac cgcgcacctg gtggagctgg ttaagcagcg cattgaggtc	6180
acggatggaa ggctacaagc ggcctttgtc gtgtcgcggg cgatcaaagg cacgcgcac	6240
ggcgggtgagg ttgccgaggc gctggccggg tacgagctgc ccattcttga gtcccgtatc	6300
acgcagcgcg tgagctaccc aggcactgcc gccgccggca caaccgttct tgaatcagaa	6360
cccgagggcg acgctgcccg cgaggctcag gcgctggccg ctgaaattaa atcaaaactc	6420
atttgagtta atgaggtaaa gagaaaatga gcaaaagcac aaacacgcta agtgccggcc	6480
gtccgagcgc acgcagcagc aaggctgcaa cgttggccag cctggcagac acgccagcca	6540
tgaagcgggt caactttcag ttgccggcgg aggatcacac caagctgaag atgtacgcgg	6600
tacgccaagg caagaccatt accgagctgc tatctgaata catcgcgag ctaccagagt	6660
<u>aa</u> atgagcaa atgaataaat gagtagatga attttagcgg ctaaaggagg cggcatggaa	6720
aatcaagaac aaccaggcac cgacgccgtg gaatgcccc tgtgtggagg aacgggcggt	6780



tggccaggcg taagcggctg ggttgtctgc cggccctgca atggcactgg aacccccaaag	6840
cccgaggaat cggcgtgacg gtcgcaaacc atccggccccg gtacaaatcg gcgcggcgct	6900
gggtgatgac ctggtggaga agttgaaggc cgcgcaggcc gccagcggc aacgcatacga	6960
ggcagaagca cggcccggtg aatcgtggca agcggccgct gatcgaatcc gcaaagaatc	7020
ccggcaaccg ccggcagccg gtgcgccgtc gattaggaag ccgccaagg gcgacgagca	7080
accagatttt ttcgttccga tgctctatga cgtgggcacc cgcgatagtc gcagcatcat	7140
ggacgtggcc gttttccgtc tgcgaagcg tgaccgacga gctggcgagg tgatccgcta	7200
cgagcttcca gacgggcacg tagaggtttc cgcagggccg gccggcatgg ccagtgtgtg	7260
ggattacgac ctggtactga tggcggtttc ccatctaacc gaatccatga accgataaccg	7320
ggaaggggaag ggagacaagc ccggccgcgt gttccgtcca cacgttgccg acgtactcaa	7380
gttctgccgg cgagccgatg gcggaaagca gaaagacgac ctggtagaaa cctgcattcg	7440
gttaaacacc acgcacgttg ccatgcagcg tacgaagaag gccagaacg gccgcctggg	7500
gacggtatcc gaggtgaag ccttgattag ccgctacaag atcgtaaaga gcgaaaccgg	7560
gcggccggag tacatcgaga tcgagctagc tgattggatg taccgcgaga tcacagaagg	7620
caagaaccgg gacgtgctga cggttcacc cgtacttt ttgatcgatc ccggcatcgg	7680
ccgttttctc taccgcctgg cacgcgcgc cgcaggcaag gcagaagcca gatggttgtt	7740
caagacgac tacgaacgca gtggcagcgc cggagagttc aagaagttct gtttcaccgt	7800
gcgcaagctg atcgggtcaa atgacctgcc ggagtacgat ttgaaggagg aggcggggca	7860
ggctggcccc atcctagtca tgcgctaccg caacctgac gagggcgaag catccgccgg	7920
ttcctaattgt acggagcaga tgctagggca aattgcccta gcaggggaaa aaggtcgaaa	7980
aggtctcttt cctgtggata gcacgtacat tgggaacca aagccgtaca ttgggaaccg	8040
gaacccttac attgggaacc caaagccgta cattgggaac cggtcacaca tgtaagtgac	8100
tgatataaaa gagaaaaaag gcgatttttc cgcctaaaac tctttaaaac ttattaaaac	8160
tcttaaaacc cgcctggcct gtgcataact gtctggccag cgcacagccg aagagctgca	8220
aaaagcgcct acccttcggt cgctgcgctc cctacgcccc gccgcttcgc gtcggcctat	8280
cgcggccgct ggccgctcaa aaatggctgg cctacggcca ggcaatctac cagggcgcgg	8340
acaagccgcg ccgtcgccac tcgaccgccg gcgcccacat caaggcacc tgcctcgcgc	8400
gtttcggtga tgacggtgaa aacctctgac acatgcagct cccggagacg gtcacagctt	8460
gtctgtaagc ggatgccggg agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg	8520
ggtgtcgggg cgcagccatg acccagtcac gtagcgatag cggagtgtat actggcttaa	8580

```

ctatgcggca tcagagcaga ttgtactgag agtgcaccat atgcggtgtg aaataccgca 8640
cagatgcgta aggagaaaaat accgcatcag gcgctcttcc gcttcctcgc tcaactgactc 8700
gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct cactcaaagg cggtaatacg 8760
gttatccaca gaatcagggg ataacgcagg aaagaacatg tgagcaaaaag gccagcaaaa 8820
ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc cataggctcc gccccctga 8880
cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aacccgacag gactataaag 8940
ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga ccctgccgct 9000
taccggatac ctgtccgcct ttctcccttc gggaaagcgtg gcgctttctc atagctcacg 9060
ctgtaggtat ctcagttcgg tgtaggctgt tcgctccaag ctgggctgtg tgcacgaacc 9120
ccccgttcag cccgaccgct gcgccttata cggtaaactat cgtcttgagt ccaaccgggt 9180
aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca gagcgaggta 9240
tgtaggcggt gctacagagt tcttgaagtg gtggcctaac tacggctaca ctagaaggac 9300
agtatttggt atctgcgctc tgctgaagcc agttaccttc ggaaaaagag ttggtagctc 9360
ttgatccggc aaacaaacca ccgctggtag cggtggtttt tttgtttgca agcagcagat 9420
tacgcgcaga aaaaaaggat ctcaagaaga tccggaaaac gcaagcgcaa agagaaagca 9480
ggtagcttgc agtgggctta catggcgata gctagactgg gcggttttat ggacagcaag 9540
cgaaccggaa ttgcc 9555

```

&lt;210&gt; 7

&lt;211&gt; 4182

&lt;212&gt; DNA

&lt;213&gt; Artificial: plasmid pRVA44

&lt;400&gt; 7

```

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatacctg caggcaattg gtacctacgt atgcatggcg cgccatatgc ccgggccctg 240
tacagcggcc gcgttcctac gcagcaggtc tcatcaagac gatctaccg agtaacaatc 300
tccaggagat caaatacctt cccaagaagg ttaaagatgc agtcaaaaga ttcaggacta 360
attgcatcaa gaacacagag aaagacatat ttctcaagat cagaagtact attccagtat 420
ggacgattca aggcttgctt cataaaccaa ggcaagtaat agagattgga gtctctaaaa 480
aggtagttcc tactgaatct aaggccatgc atggagtcta agattcaaat cgaggatcta 540
acagaactcg ccgtgaagac tggcgaacag ttcatacaga gtcttttacg actcaatgac 600

```

aagaagaaaa	tcttcgtcaa	catggtggag	cacgacactc	tggtctactc	caaaaatgtc	660
aaagatacag	tctcagaaga	ccaaagggct	attgagactt	ttcaacaaag	gataatttcg	720
ggaaacctcc	tgggattcca	ttgcccagct	atctgtcact	tcatcgaaag	gacagtagaa	780
aaggaaggtg	gtcctacaa	atgccatcat	tgcgataaag	gaaaggctat	cattcaagat	840
gcctctgccg	acagtgggtc	caaagatgga	ccccaccca	cgaggagcat	cgtggaaaaa	900
gaagacgttc	caaccacgtc	ttcaaagcaa	gtggattgat	gtgacatctc	cactgacgta	960
agggatgacg	cacaatccca	ctatccttcg	caagaccctt	cctctatata	aggaagttca	1020
tttcatttgg	agaggacacg	ctgaaatcac	cagtctctct	ctataaatct	atctctctct	1080
ctataaccat	ggacccagaa	cgacgcccg	ccgacatccg	ccgtgccacc	gaggcggaca	1140
tgccggcggg	ctgcaccatc	gtcaaccact	acatcgagac	aagcacggtc	aacttccgta	1200
ccgagccgca	ggaaccgcag	gagtggacgg	acgacctcgt	ccgtctgcgg	gagcgctatc	1260
cctggctcgt	cgccgaggtg	gacggcgagg	tcgccggcat	cgcctacgcg	ggcccctgga	1320
aggcacgcaa	cgcctacgac	tggacggccg	agtcgaccgt	gtacgtctcc	ccccgccacc	1380
agcggacggg	actgggctcc	acgctctaca	cccacctgct	gaagtccttg	gaggcacagg	1440
gcttcaagag	cgtggtoget	gtcatcgggc	tgcccaacga	cccgagcgtg	cgcattgcacg	1500
aggcgctcgg	atatgcccc	cgcggcatgc	tgccggcggc	cggcttcaag	cacgggaact	1560
ggcatgacgt	gggtttctgg	cagctggact	tcagcctgcc	ggtaccgccc	cgtccgggtcc	1620
tgcccgtcac	cgagatctga	tctcacgcgt	ctaggatccg	aagcagatcg	ttcaaacatt	1680
tggcaataaa	gtttcttaag	attgaatcct	gttgccggtc	ttgcgatgat	tatcatataa	1740
tttctgttga	attacgttaa	gcatgtaata	attaacatgt	aatgcatgac	gttattttatg	1800
agatgggttt	ttatgattag	agtcccgcaa	ttatacattt	aatacgcgat	agaaaacaaa	1860
atatagcgcg	caaactagga	taaattatcg	cgcgcgggtg	catctatggt	actagatcgg	1920
gaagatcctc	tagagcgatc	gcaagcttgg	cgtaatcatg	gtcatagctg	tttctgtgt	1980
gaaattgtta	tccgctcaca	attccacaca	acatacgagc	cggaagcata	aagtgtaaag	2040
cctgggggtgc	ctaattgagt	agctaactca	cattaattgc	gttgcgctca	ctgcccgttt	2100
tccagtcggg	aaacctgtcg	tgccagctgc	attaatgaat	cggccaacgc	gcgggggagag	2160
gcggtttgcg	tattggggcg	tcttccgctt	cctcgctcac	tgactcgctg	cgtcgggtcg	2220
ttcggctgcg	gcgagcggta	tcagctcact	caaaggcggg	aatacggtta	tccacagaat	2280
caggggataa	cgcaggaaag	aacatgtgag	caaaaggcca	gcaaaaggcc	aggaaccgta	2340
aaaaggccgc	gttgctggcg	tttttccata	ggctccgccc	ccctgacgag	catcacaaaa	2400
atcgacgctc	aagtcagagg	tggcgaaacc	cgacaggact	ataaagatac	caggcgtttc	2460

```

ccccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 2520
ccgcctttct cccttcggga agcgtggcgc tttctcaaag ctcacgctgt aggtatctca 2580
gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg 2640
accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccggtgaaga cacgacttat 2700
cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 2760
cagagttctt gaagtggtag cctaactacg gctacactag aagaacagta tttggtatct 2820
gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 2880
aaaccaccgc tggtagcggg ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 2940
aaggatctca agaagatcct ttgatctttt ctacggggtc tgacgctcag tggaacgaaa 3000
actcacgtta agggattttg gtcattgagat tatcaaaaag gatcttcacc tagatccttt 3060
taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tgggtctgaca 3120
gttaccaatg cttaatcagt gaggcacctt tctcagcgat ctgtctattt cgttcatcca 3180
tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggctta ccatctggcc 3240
ccagtgtctg aatgataccg cgagaccac gctcaccggc tccagattta tcagcaataa 3300
accagccagc cggaagggcc gagcgcagaa gtggtcctgc aactttatcc gcctccatcc 3360
agtctattaa ttgttgccgg gaagctagag taagtagttc gccagttaat agtttgcgca 3420
acgttggtgc cattgctaca ggcacgtggt tgtcacgctc gtcgtttggg atggcttcat 3480
tcagctccgg ttcccaacga tcaaggcgag ttacatgatc ccccatgttg tgcaaaaaag 3540
cggttagctc cttcggctct ccgatcgttg tcagaagtaa gttggccgca gtgttatcac 3600
tcattggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 3660
ctgtgactgg tgagtactca accaagtcatt tctgagaata gtgtatgcgg cgaccgagtt 3720
gctcttgccc ggcgtcaata cgggataata ccgcgccaca tagcagaact ttaaaagtgc 3780
tcatcattgg aaaacgttct tcggggcgaa aactctcaag gatcttaccg ctggtgagat 3840
ccagttcgat gtaaccact cgtgcacca actgatcttc agcatctttt actttcacca 3900
gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 3960
cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg 4020
gttattgtct catgagcgga tacatatctg aatgtattta gaaaaataaa caaatagggg 4080
ttccgcgcac atttccccga aaagtgccac ctgacgtcta agaaaccatt attatcatga 4140
cattaacctt taaaaatagg cgtatcacga ggccctttcg tc 4182

```

<211> 16  
 <212> DNA  
 <213> Artificial: primer MDB327

<220>  
 <221> misc\_feature  
 <222> (1)..(16)  
 <223> "n" = a,c,t or g; "s" = c or g; "w" = a or t

<400> 8  
 ntgaggwtcn wgtsat 16

<210> 9  
 <211> 22  
 <212> DNA  
 <213> Artificial: primer MLD015

<400> 9  
 tggttcctag cgtgagccag tg 22

<210> 10  
 <211> 21  
 <212> DNA  
 <213> Artificial: primer MLD016

<400> 10  
 agctgctgct cttgcctctg t 21

<210> 11  
 <211> 16  
 <212> DNA  
 <213> Artificial: primer MDB612

<220>  
 <221> misc\_feature  
 <222> (1)..(16)  
 <223> "n" = a,c,t or g; "s" = c or g; "w" = a or t

<400> 11  
 ngtgctswga nawgat 16

<210> 12  
 <211> 22  
 <212> DNA  
 <213> Artificial: primer MDB053

<400> 12  
 catgacgtgg gtttctggca gc 22

<210> 13  
 <211> 21  
 <212> DNA  
 <213> Artificial: primer MDB356

<400> 13  
 aatcctgttg ccggtcttgc g 21

<210> 14  
 <211> 22  
 <212> DNA  
 <213> Artificial: primer DPA017

<400> 14  
 gattagagtc ccgcaattat ac 22

<210> 15  
 <211> 20  
 <212> DNA  
 <213> Artificial: primer MLD019

<400> 15  
 caagatgcga gcaactatgt 20

<210> 16  
 <211> 200  
 <212> DNA  
 <213> Artificial: sequence comprising target site deletion

<400> 16  
 tcttggacca caattttgag cttttcattt tctattttga ggataatagt ttattccctc 60  
 aaggaactat tcaactgagc ttaatatctc aatttttttt aacatatgac tataagtatc 120  
 ctccaaatat ttaaaaagaa taccaccatt atccgaatct tctttaaaat ctgttagaac 180  
 acggtttgga atagtggtag 200

<210> 17  
 <211> 21  
 <212> DNA  
 <213> Artificial: primer GHI01

<400> 17  
 aacctaggct gctgaaggag c 21

<210> 18  
 <211> 21  
 <212> DNA  
 <213> Artificial: primer GHI02

<400> 18  
 caactcctcc agtcattctcc g 21